



# *Bolldex*<sup>®</sup> & *Cryptex*<sup>®</sup>

Virus based biocontrol products for the effective control of key lepidoptera pests in Stone Fruit

*Bolldex*<sup>®</sup> and *Cryptex*<sup>®</sup> are virus based biocontrol products produced by world renowned virus specialists, Andermatt Biocontrol in Switzerland.

Virus based biocontrol products:

- are extremely host specific, with zero negative effects on non-target organisms, ideal for use in IPM programs.
- can be applied during flowering as they have no negative effects on bees
- are residue free with no MRL constraints (ie. ideal for use on organic and export crops)
- have a completely different mode of action to conventional chemistry, making them ideal IPM partners from a resistance management perspective

## *Bolldex*<sup>®</sup>

(Reg. no. L 8895, Act 36 of 1947)



*Bolldex*<sup>®</sup> offers targeted control of Africa Bollworm.

## *Cryptex*<sup>®</sup>

(Reg. no. L 8037, Act 36 of 1947)



*Cryptex*<sup>®</sup> specifically controls False Codling Moth.

### Application:

- *Bolldex*<sup>®</sup> is registered on all crops at a standard application rate of 200ml/ha. On Stone fruit applications should be made during flowering or as soon as Bollworm eggs are observed.
- *Cryptex*<sup>®</sup> should be applied at 60ml/ha 10-14 days after the 1st moth peak. Applications should be repeated at 35ml/ha every 10-14 days thereafter.
- *Cryptex*<sup>®</sup> and *Bolldex*<sup>®</sup> can be mixed with most other agricultural inputs provided the pH of the tank solution is between 5-8. Copper is the one exception. Tank mixes with Copper should be avoided.
- For detailed application instructions please refer to the registered label.



Madumbi Sustainable Agriculture (Pty) Ltd  
+27 (0) 33 342 3984 | support@madumbi.co.za | [www.madumbi.co.za](http://www.madumbi.co.za)

Backed by Science. Loved by Nature.